

**SRI LANKA STANDARD 1350 : 2016**

**ISO 22717 : 2015**

**UDC 665.58:579.63**

**METHOD OF TEST FOR  
THE DETECTION OF *Pseudomonas aeruginosa* IN  
COSMETICS  
(First Revision)**

**SRI LANKA STANDARDS INSTITUTION**



**Sri Lanka Standard**  
**METHOD OF TEST FOR THE DETECTION OF**  
***Pseudomonas aeruginosa* IN COSMETICS**  
**(First Revision)**

**SLS 1350 : 2016**  
**ISO 22717 : 2015**

**Gr. G**

*Copyright Reserved*  
**SRI LANKA STANDARDS INSTITUTION**  
**17, Victoria Place**  
**Elvitigala Mawatha**  
**Colombo 08**  
**SRI LANKA**

Sri Lanka Standards are subject to periodical revision in order to accommodate the progress made by industry. Suggestions for improvement will be recorded and brought to the notice of the Committees to which the revisions are entrusted.

This standard does not purport to include all the necessary provisions of a contract.

© **SLSI 2016**

All right reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the SLSI.

**Sri Lanka Standard**  
**METHOD OF TEST FOR THE DETECTION OF**  
***Pseudomonas aeruginosa* IN COSMETICS**  
**(First Revision)**

**FOREWORD**

This Sri Lanka Standard was approved by the Sectoral Committee on Chemical and Polymer Technology and was authorized for adoption and publication as a Sri Lanka Standard by the Council of the Sri Lanka Standards Institution on 2016-10-27.

This Sri Lanka Standard was first published in 2008 which was an adoption of ISO 22717 : 2006 Cosmetics – Microbiology – Detection of *Pseudomonas aeruginosa*. The International Standard ISO 22717 : 2006 has been technically revised in 2015. ISO 22717 : 2015 which gives general guidelines for the detection and identification of *Pseudomonas aeruginosa* in cosmetic products has been accepted to adopt as the first revision to **SLS 1350 : 2016**

This Standard is identical with ISO 22717 : 2015 Cosmetics – Microbiology – Detection of *Pseudomonas aeruginosa*, published by the International Organization for Standardization (ISO).

**TERMINOLOGY AND CONVENTIONS :**

The text of the International Standard has been accepted as suitable for publication, without deviation, as a Sri Lanka Standard. However, certain terminology and conventions are not identical with those used in Sri Lanka Standards. Attention is therefore drawn to the following :

- a) Wherever the words ‘International Standard’ appear referring to a particular standard, they should be interpreted as “Sri Lanka Standard”.
- b) The comma has been used throughout as a decimal marker. In Sri Lanka Standards it is the current practice to use the full point at the base as the decimal marker.
- c) Wherever page numbers are quoted, they are ISO page numbers.

SLS 1350 : 2016  
ISO 22717 : 2015

## **Cross References**

### **International Standard**

ISO 21148 Cosmetics – Microbiology –  
General instructions for microbiological  
examination

EN 12353 Chemical disinfectants and  
antiseptics – Preservation of microbial stains  
used for the determination of bactericidal and  
fungicidal activity

### **Corresponding Sri Lanka Standard**

No corresponding Sri Lanka Standard

No corresponding Sri Lanka Standard

INTERNATIONAL  
STANDARD

SLS 1350:2016

**ISO**  
**22717**

Second edition  
2015-11-15

---

---

**Cosmetics — Microbiology —  
Detection of *Pseudomonas aeruginosa***

*Cosmétiques — Microbiologie — Détection de Pseudomonas  
aeruginosa*



Reference number  
ISO 22717:2015(E)

© ISO 2015



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2015, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Ch. de Blandonnet 8 • CP 401  
CH-1214 Vernier, Geneva, Switzerland  
Tel. +41 22 749 01 11  
Fax +41 22 749 09 47  
copyright@iso.org  
www.iso.org



# Contents

	Page
Foreword .....	iv
Introduction .....	v
<b>1 Scope</b> .....	<b>1</b>
<b>2 Normative references</b> .....	<b>1</b>
<b>3 Terms and definitions</b> .....	<b>1</b>
<b>4 Principle</b> .....	<b>2</b>
<b>5 Diluents and culture media</b> .....	<b>2</b>
5.1 General .....	2
5.2 Diluent for the bacterial suspension (tryptone sodium chloride solution) .....	3
5.2.1 General .....	3
5.2.2 Composition .....	3
5.2.3 Preparation .....	3
5.3 Culture media .....	3
5.3.1 General .....	3
5.3.2 Agar medium for the suitability test (see Clause 11) [soybean–casein digest agar medium (SCDA) or tryptic soy agar (TSA)] .....	3
5.3.3 Enrichment broth .....	4
5.3.4 Selective agar medium for isolation of <i>Pseudomonas aeruginosa</i> .....	5
5.3.5 Selective agar medium for confirmation of <i>Pseudomonas aeruginosa</i> .....	5
<b>6 Apparatus and glassware</b> .....	<b>6</b>
<b>7 Strains of microorganisms</b> .....	<b>6</b>
<b>8 Handling of cosmetic products and laboratory samples</b> .....	<b>6</b>
<b>9 Procedure</b> .....	<b>6</b>
9.1 General recommendation .....	6
9.2 Preparation of the initial suspension in the enrichment broth .....	6
9.2.1 General .....	6
9.2.2 Water-miscible products .....	7
9.2.3 Water-immiscible products .....	7
9.2.4 Filterable products .....	7
9.3 Incubation of the inoculated enrichment broth .....	7
9.4 Detection and Identification of <i>Pseudomonas aeruginosa</i> .....	7
9.4.1 Isolation .....	7
9.4.2 Identification of <i>Pseudomonas aeruginosa</i> .....	7
<b>10 Expression of results (detection of <i>Pseudomonas aeruginosa</i>)</b> .....	<b>8</b>
<b>11 Neutralization of the antimicrobial properties of the product</b> .....	<b>8</b>
11.1 General .....	8
11.2 Preparation of the inoculum .....	8
11.3 Suitability of the detection method .....	8
11.3.1 Procedure .....	8
11.3.2 Interpretation of suitability test results .....	9
<b>12 Test report</b> .....	<b>9</b>
<b>Annex A (informative) Other enrichment broths</b> .....	<b>10</b>
<b>Annex B (informative) Neutralizers of antimicrobial activity of preservatives and rinsing liquids</b> .....	<b>12</b>
<b>Bibliography</b> .....	<b>13</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: [Foreword - Supplementary information](#)

The committee responsible for this document is ISO/TC 217, *Cosmetics*.

This second edition cancels and replaces the first edition (ISO 22717:2006), of which it constitutes a minor revision.

## Introduction

Microbiological examinations of cosmetic products are carried out according to an appropriate microbiological risk analysis in order to ensure their quality and safety for consumers.

Microbiological risk analysis depends on several parameters such as the following:

- potential alteration of cosmetic products;
- pathogenicity of microorganisms;
- site of application of the cosmetic product (hair, skin, eyes, mucous membranes);
- type of users (adults, children under 3 years).

For cosmetics and other topical products, the detection of skin pathogens such as *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans* may be relevant because they can cause skin or eye infections. The detection of other kinds of microorganism might be of interest since these microorganisms (including indicators of faecal contamination e.g. *Escherichia coli*) suggest hygienic failure during the manufacturing process.



# Cosmetics — Microbiology — Detection of *Pseudomonas aeruginosa*

## 1 Scope

This International Standard gives general guidelines for the detection and identification of the specified microorganism *Pseudomonas aeruginosa* in cosmetic products. Microorganisms considered as specified in this International Standard might differ from country to country according to national practices or regulations.

In order to ensure product quality and safety for consumers, it is advisable to perform an appropriate microbiological risk analysis to determine the types of cosmetic product to which this International Standard is applicable. Products considered to present a low microbiological (see ISO 29621) risk include those with low water activity, hydro-alcoholic products, extreme pH values, etc.

The method described in this International Standard is based on the detection of *Pseudomonas aeruginosa* in a non-selective liquid medium (enrichment broth), followed by isolation on a selective agar medium. Other methods may be appropriate, depending on the level of detection required.

**NOTE** For the detection of *Pseudomonas aeruginosa*, subcultures can be performed on non-selective culture media followed by suitable identification steps (e.g. using identification kits).

Because of the large variety of cosmetic products within this field of application, this method may not be appropriate in every detail for some products (e.g. certain water immiscible products). Other International Standards (ISO 18415) may be appropriate. Other methods (e.g. automated) may be substituted for the tests presented here provided that their equivalence has been demonstrated or the method has been otherwise shown to be suitable.

## 2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 21148:2005, *Cosmetics — Microbiology — General instructions for microbiological examination*

EN 12353, *Chemical disinfectants and antiseptics — Preservation of test organisms used for the determination of bactericidal (including Legionella), mycobactericidal, sporicidal, fungicidal and virucidal (including bacteriophages) activity*

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

### 3.1

#### **product**

portion of an identified cosmetic product received in the laboratory for testing

### 3.2

#### **sample**

portion of the product (at least 1 g or 1 ml) that is used in the test to prepare the initial suspension

**3.3**  
**initial suspension**

suspension (or solution) of the sample in a defined volume of an appropriate enrichment broth

**3.4**  
**sample dilution**

dilution of the initial suspension

**3.5**  
**specified microorganism**

aerobic mesophilic bacterium or yeast that is undesirable in a cosmetic product and is recognized as a skin pathogen species that may be harmful for human health or as an indication of hygienic failure in the manufacturing process

**3.6**  
***Pseudomonas aeruginosa***

gram-negative rod, motile; smooth colonies pigmented brown or greenish

Note 1 to entry: The main characteristics for identification are: growth on selective cetrimide agar medium, oxidase positive, production of diffusible fluorescent pigments and production of a soluble phenazine pigment (pyocyanin) in suitable media.

Note 2 to entry: *Pseudomonas aeruginosa* may be isolated from a wide variety of environmental sources, especially in water and has a very high potential to spoil many different substrates. It may produce infections of human skin or eye area. It is undesirable in cosmetic products for its potential pathogenicity and its capacity to affect the physico-chemical properties of the cosmetic formula.

**3.7**  
**enrichment broth**

non-selective liquid medium containing suitable neutralizers and/or dispersing agents and demonstrated to be suitable for the product under test

## **4 Principle**

The first step of the procedure is to perform an enrichment by using a non-selective broth medium to increase the number of microorganisms without the risk of inhibition by the selective ingredients that are present in selective/differential growth media.

The second step of the test (isolation) is performed on a selective medium followed by identification tests.

The possible inhibition of microbial growth by the sample shall be neutralized to allow the detection of viable microorganisms.<sup>[1]</sup> In all cases and whatever the methodology, the neutralization of the antimicrobial properties of the product shall be checked and demonstrated (see [Clause 11](#)).

## **5 Diluents and culture media**

### **5.1 General**

General instructions are given in ISO 21148. When water is mentioned in this International Standard, use distilled water or purified water as specified in ISO 21148.

The enrichment broth is used to disperse the sample and to increase the initial microbial population. It may contain neutralizers if the specimen to be tested has antimicrobial properties. The efficacy of the neutralization shall be demonstrated (see [Clause 11](#)). Information relative to suitable neutralizers is given in [Annex B](#).

The enrichment broth ([5.3.3.1](#)), or any of the ones listed in [Annex A](#), is suitable for checking the presence of *Pseudomonas aeruginosa* in accordance with this International Standard provided that it has been demonstrated to be suitable in accordance with [Clause 11](#).

Other diluents and culture media may be used if it has been demonstrated that they are suitable for use.

## 5.2 Diluent for the bacterial suspension (tryptone sodium chloride solution)

### 5.2.1 General

The diluent is used for the preparation of bacterial suspension used for the suitability test procedure (see [Clause 11](#)).

### 5.2.2 Composition

— tryptone, pancreatic digest of casein	1,0 g
— sodium chloride	8,5 g
— water	1 000 ml

### 5.2.3 Preparation

Dissolve the components in water by mixing while heating. Dispense into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,0 \pm 0,2$  when measured at room temperature.

## 5.3 Culture media

### 5.3.1 General

Culture media may be prepared using the descriptions provided below or from dehydrated culture media according to the instructions of the manufacturer. The instructions provided by the supplier of the media should be followed.

NOTE Ready-to-use media can be used when their composition and/or growth yields are comparable to those of the formulae given herein.

### 5.3.2 Agar medium for the suitability test (see [Clause 11](#)) [soybean–casein digest agar medium (SCDA) or tryptic soy agar (TSA)]

#### 5.3.2.1 Composition

— pancreatic digest of casein	15,0 g
— papaic digest of soybean meal	5,0 g
— sodium chloride	5,0 g
— agar	15,0 g
— water	1 000 ml

#### 5.3.2.2 Preparation

Dissolve the components or the dehydrated complete medium in the water by mixing while heating. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,3 \pm 0,2$  when measured at room temperature.

### 5.3.3 Enrichment broth

#### 5.3.3.1 Eugon LT 100 broth

##### 5.3.3.1.1 General

This medium contains ingredients that neutralize inhibitory substances present in the sample: lecithin and polysorbate 80, and dispersing agent: octoxynol 9.

##### 5.3.3.1.2 Composition

— pancreatic digest of casein	15,0 g
— papaic digest of soybean meal	5,0 g
— L-cystine	0,7 g
— sodium chloride	4,0 g
— sodium sulfite	0,2 g
— glucose	5,5 g
— egg lecithin	1,0 g
— polysorbate 80	5,0 g
— octoxynol 9	1,0 g
— water	1 000 ml

##### 5.3.3.1.3 Preparation

Dissolve the components polysorbate 80, octoxynol 9 and egg lecithin successively into boiling water until their complete dissolution. Dissolve the other components by mixing while heating. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,0 \pm 0,2$  when measured at room temperature.

#### 5.3.3.2 Other enrichment broths

Other enrichment broths may be used as appropriate (see [Annex A](#)).



### 5.3.4 Selective agar medium for isolation of *Pseudomonas aeruginosa*

#### 5.3.4.1 Cetrimide agar medium

##### 5.3.4.1.1 Composition

— pancreatic digest of gelatin	20,0 g
— magnesium chloride	1,4 g
— potassium sulfate	10,0 g
— cetrimide (cetyltrimethylammonium bromide)	0,3 g
— agar	13,6 g
— glycerol	10,0 ml
— water	1 000 ml

##### 5.3.4.1.2 Preparation

Dissolve all solid components in the water and add the glycerol. Heat, with frequent agitation, and boil for 1 min to effect dissolution.

Dispense in suitable flasks and sterilize at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,2 \pm 0,2$  when measured at room temperature.

### 5.3.5 Selective agar medium for confirmation of *Pseudomonas aeruginosa*

#### 5.3.5.1 *Pseudomonas* agar medium for detection of pyocyanin (*Pseudomonas* agar P)

##### 5.3.5.1.1 Composition

— pancreatic digest of gelatin	20,0 g
— anhydrous magnesium chloride	1,4 g
— anhydrous potassium sulfate	10,0 g
— agar	15,0 g
— glycerol	10,0 ml
— water	1 000 ml

##### 5.3.5.1.2 Preparation

Dissolve all solid components in the water, and add the glycerol. Heat, with frequent agitation, and boil for 1 min to effect dissolution.

Dispense in suitable flasks and sterilize at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,2 \pm 0,2$  when measured at room temperature.

## 6 Apparatus and glassware

Use the laboratory equipment, apparatus and glassware as described in ISO 21148.

## 7 Strains of microorganisms

For the verification of the suitability test conditions, the following representative strain is used:

*Pseudomonas aeruginosa* ATCC<sup>1)</sup> 9027 (equivalent strain: CIP<sup>2)</sup> 82118 or NCIMB<sup>3)</sup> 8626 or NBRC<sup>4)</sup> 13275 or KCTC<sup>5)</sup> 2513 or other equivalent national collection strain).

The culture should be reconstituted according to the procedures provided by the supplier of reference strain.

The strain may be stored in the laboratory in accordance with EN 12353.

## 8 Handling of cosmetic products and laboratory samples

If necessary, store products to be tested at room temperature.

Do not incubate, refrigerate or freeze products and samples before or after analysis.

Sampling of cosmetic products to be analysed should be carried out as described in ISO 21148. Analyse samples as described in ISO 21148 and according to the procedure in [Clause 9](#).

## 9 Procedure

### 9.1 General recommendation

Use sterile material, equipment and aseptic techniques to prepare the sample, initial suspension and dilutions. In the case of the preparation of the initial suspension in an appropriate solubilizing agent, the time which elapses between the end of preparation and the moment the inoculum comes into contact with the enrichment broth shall not exceed 45 min, unless specifically mentioned in the established protocols or documents.

### 9.2 Preparation of the initial suspension in the enrichment broth

#### 9.2.1 General

The enrichment is prepared from a sample ([3.2](#)) of at least 1 g or 1 ml of the well-mixed product under test, which is dispersed in at least 9 ml of enrichment broth.

Note S, the exact weight or volume of the sample.

The method shall be checked to ensure that the composition (neutralizer eventually added) and the volume of the broth perform satisfactorily (see [11.3](#)).

NOTE In some cases, and when possible, filtration of the cosmetic product through a membrane that is afterwards immersed in the enrichment broth, facilitates the neutralization of the antimicrobial properties of the product (see [11.3](#)).

- 
- 1) ATCC = American Type Culture Collection.
  - 2) CIP = Institut Pasteur Collection.
  - 3) NCIMB = National Collection of Industrial and Marine Bacteria.
  - 4) NBRC = National Biological Resource center.
  - 5) KCTC = Korean Collection for type culture.

### 9.2.2 Water-miscible products

Transfer the sample, *S*, of product to a suitable container containing an appropriate volume of broth.

### 9.2.3 Water-immiscible products

Transfer the sample, *S*, of product to a suitable container containing a suitable quantity of solubilizing agent (*e.g.* Polysorbate 80).

Disperse the sample within the solubilizing agent and add an appropriate volume of broth.

### 9.2.4 Filterable products

Use a membrane filter having a nominal pore size of not greater than 0,45 µm.

Transfer the sample, *S*, on to the membrane in a filtration apparatus (see ISO 21148). Filter immediately and wash the membrane using defined volumes of water and/or diluent.

Transfer and immerse the membrane into a tube or flask of suitable size containing an appropriate volume of broth.

## 9.3 Incubation of the inoculated enrichment broth

Incubate the initial suspension prepared in broth (see 9.2) at 32,5 °C ± 2,5 °C for at least 20 h (maximum 72 h).

## 9.4 Detection and Identification of *Pseudomonas aeruginosa*

### 9.4.1 Isolation

Using a sterile loop, streak an aliquot of the incubated enrichment broth on the surface of cetrimide agar medium in order to obtain isolated colonies.

Invert the Petri dish and then incubate at 32,5 °C ± 2,5 °C for at least 24 h (maximum 48 h).

Check for characteristic colonies (see [Table 1](#)).

**Table 1 — Morphological characteristics of *Pseudomonas aeruginosa* on selective medium**

Selective medium	Characteristic colonial morphology of <i>Pseudomonas aeruginosa</i>
Cetrimide agar medium	Yellow-green pigment (pyocyanin), which fluoresces under UV light.

### 9.4.2 Identification of *Pseudomonas aeruginosa*

#### 9.4.2.1 General

Proceed to the following tests, for the suspect colonies isolated on the cetrimide agar medium. The presence of *Pseudomonas aeruginosa* may be confirmed by other suitable, cultural and biochemical tests.

#### 9.4.2.2 Gram's stain

This test is described in ISO 21148.

Check for Gram-negative rods.

#### 9.4.2.3 Oxidase test

This test is described in ISO 21148.

Check for oxidase positive test.

#### 9.4.2.4 Culture on *Pseudomonas* agar medium for detection of pyocyanin

Inoculate the surface of the *Pseudomonas* agar medium for detection of pyocyanin with suspect isolated colonies grown on cetrimide agar medium, so that individual colonies develop. Incubate at  $32,5\text{ °C} \pm 2,5\text{ °C}$ .

Check for bacterial growth after 24 h, 48 h and 72 h. *Pseudomonas aeruginosa* forms colonies surrounded by a blue to green zone due to pyocyanin formation or with a red to dark brown zone due to pyorubin production.

### 10 Expression of results (detection of *Pseudomonas aeruginosa*)

If the identification of the colonies confirms the presence of this species, express the result as:

— presence of *Pseudomonas aeruginosa* in the sample, *S*.

If no growth after enrichment is observed and/or if the identification of the colonies does not confirm the presence of this species, express the result as:

— absence of *Pseudomonas aeruginosa* in the sample, *S*.

### 11 Neutralization of the antimicrobial properties of the product

#### 11.1 General

The different tests described below demonstrate that the microorganism can grow in analysis conditions.

#### 11.2 Preparation of the inoculum

Prior to the test, inoculate the surface of soybean casein digest agar (SCDA) or other suitable (non-selective, non-neutralizing) medium with *Pseudomonas aeruginosa*. Incubate the plate at  $32,5\text{ °C} \pm 2,5\text{ °C}$  for 18 h to 24 h.

To harvest the culture use a sterile loop, streak the surface of the culture and re-suspend in the diluent to obtain a calibrated suspension of about  $1 \times 10^8$  CFU per ml (e.g. using spectrophotometer, ISO 21148:2005, Annex C).

Use this calibrated suspension and its dilutions within 2 h.

#### 11.3 Suitability of the detection method

##### 11.3.1 Procedure

**11.3.1.1** In tubes of 9 ml of diluent, prepare a dilution of the calibrated suspension in order to obtain a final count between 100 CFU and 500 CFU per ml. To count the final concentration of viable microorganisms in the diluted calibrated suspension, transfer 1 ml of the suspension into a Petri dish and pour on 15 ml to 20 ml of the melted agar medium kept in a water bath at no more than 48°C. Let solidify and then incubate at  $32,5\text{ °C} \pm 2,5\text{ °C}$  for 20 h to 24 h.

**11.3.1.2** Prepare in duplicate, the initial suspension in the conditions chosen for the test (at least 1 g or 1 ml of product under test, defined volume of enrichment broth) in a tube or flask. When using the membrane filtration method filter in duplicate at least 1 ml of product under test and transfer each membrane into a tube or flask containing the enrichment broth in the conditions chosen for the test.

**11.3.1.3** Introduce aseptically, 0,1 ml of the diluted calibrated suspension ([11.3.1.1](#)) of microorganisms into one tube or flask (suitability test). Mix, then incubate both tubes or flasks (suitability test and non-inoculated control) at  $32,5\text{ °C} \pm 2,5\text{ °C}$  for 20 h to 24 h.

**11.3.1.4** Perform an isolation for each tube or flask (suitability test and non-inoculated control). Using a sterile loop, streak an aliquot (same conditions as in the test) of the incubated mixture on to the surface of a Petri dish (diameter 85 mm to 100 mm) containing approximately 15 ml to 20 ml of cetrimide agar medium. Incubate the plates at  $32,5\text{ °C} \pm 2,5\text{ °C}$  for 24 h to 48 h.

### **11.3.2 Interpretation of suitability test results**

Check that the diluted calibrated suspension ([11.3.1.1](#)) of bacteria contains between 100 CFU and 500 CFU per ml.

The neutralization is verified and the detection method is satisfactory if a growth characteristic of *Pseudomonas aeruginosa* occurs on the suitability test plate and no growth occurs on the control plate.

When growth is detected on the control plate (contaminated products), the neutralization is verified and the detection method is satisfactory if *Pseudomonas aeruginosa* is recovered on the suitability test plate.

Failure of growth on the suitability test plates indicates that antimicrobial activity is still present and necessitates a modification of the conditions of the method by an increase in the volume of nutrient broth, the quantity of product remaining the same, or by incorporation of a sufficient quantity of inactivating agent in the enrichment broth, or by an appropriate combination of modifications so as to permit the growth of *Pseudomonas aeruginosa*.

If, in spite of the incorporation of suitable inactivating agents and a substantial increase in the volume of broth, it is still not possible to recover viable cultures as described above, indicate that the article is not likely to be contaminated with *Pseudomonas aeruginosa*.

## **12 Test report**

The test report shall contain the following information:

- a) a reference to this International Standard, i.e. ISO 22717:2015;
- b) all information necessary for the complete identification of the product;
- c) the method used;
- d) the results obtained;
- e) all operating details for the preparation of the initial suspension;
- f) the description of the method with the neutralizers and media used;
- g) the demonstration of the suitability of the method, even if the test has been performed separately;
- h) any point not specified in this International Standard, or regarded as optional, together with details of any incidents which may have influenced the results.

## Annex A (informative)

### Other enrichment broths

#### A.1 Soybean-casein-digest-lecithin-polysorbate 80 medium (SCDLP 80 broth)

##### A.1.1 Composition

— casein peptone	17,0 g
— soybean peptone	3,0 g
— sodium chloride	5,0 g
— potassium hydrogen phosphate	2,5 g
— glucose	2,5 g
— lecithin	1,0 g
— polysorbate 80	7,0 g
— water	1 000 ml

##### A.1.2 Preparation

Dissolve all of these components (or dehydrated complete medium) one after another in boiling water until their complete dissolution. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,2 \pm 0,2$  when measured at room temperature.

#### A.2 D/E neutralizing broth (Dey/Engley neutralizing broth)<sup>[Z]</sup>

##### A.2.1 Composition

— glucose	10,0 g
— soybean lecithin	7,0 g
— sodium thiosulfate pentahydrate	6,0 g
— polysorbate 80	5,0 g
— pancreatic digest of casein	5,0 g
— sodium bisulfite	2,5 g
— yeast extract	2,5 g
— sodium thioglycolate	1,0 g

— bromocresol purple	0,02 g
— water	1 000 ml

### A.2.2 Preparation

Dissolve all of these components (or dehydrated complete medium) one after another in boiling water until their complete dissolution. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,6 \pm 0,2$  when measured at room temperature.

## A.3 Modified letheen broth

### A.3.1 Composition

— peptic digest of meat	20,0 g
— pancreatic digest of casein	5,0 g
— beef extract	5,0 g
— yeast extract	2,0 g
— lecithin	0,7 g
— polysorbate 80	5,0 g
— sodium chloride	5,0 g
— sodium bisulfite	0,1 g
— water	1 000 ml

### A.3.2 Preparation

Dissolve successively in boiling water: polysorbate 80 and lecithin until their complete dissolution. Dissolve the other components by mixing while heating. Mix gently to avoid foam. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to  $7,2 \pm 0,2$  when measured at room temperature.

## Annex B (informative)

### Neutralizers of antimicrobial activity of preservatives and rinsing liquids

Preservative	Chemical compounds able to neutralize preservative's antimicrobial activity	Examples of suitable neutralizers and of rinse liquids (for membrane filtration methods)
Phenolic compounds: parabens, phenoxyethanol, phenylethanol, etc. anilides	Lecithin, polysorbate 80, ethylene oxide condensate of fatty alcohol, non-ionic surfactants	Polysorbate 80, 30 g/l + lecithin, 3 g/l. Ethylene oxide condensate of fatty alcohol, 7 g/l + lecithin, 20 g/l + polysorbate 80, 4 g/l. D/E neutralizing broth <sup>a</sup> Rinse liquid: distilled water; tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Quaternary ammonium compounds, cationic surfactants	Lecithin, saponin, polysorbate 80, Sodium dodecyl sulfate, Ethylene oxide condensate of fatty alcohol	Polysorbate 80, 30 g/l + sodium dodecyl sulfate, 4 g/l + lecithin, 3 g/l. Polysorbate 80, 30 g/l + saponin, 30 g/l + lecithin, 3 g/l. D/E neutralizing broth <sup>a</sup> Rinse liquid: distilled water; tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Aldehydes, formaldehyde-release agents	Glycine, histidine	Lecithin, 3 g/l + polysorbate 80, 30 g/l + L-histidine, 1 g/l. Polysorbate 80, 30 g/l + saponin, 30 g/l + L-histidine, 1 g/l + L-cysteine, 1 g/l. D/E neutralizing broth <sup>a</sup> Rinse liquid: polysorbate 80, 3 g/l + L-histidine, 0,5 g/l.
Oxidizing compounds	Sodium thiosulfate	Sodium thiosulfate, 5 g/l. Rinse liquid: sodium thiosulfate, 3 g/l.
Isothiazolinones, imidazoles	Lecithin, Saponin, amines, sulfates, mercaptans, sodium bisulfite, sodium thioglycollate	Polysorbate 80, 30 g/l + saponin, 30 g/l + lecithin, 3 g/l. Rinse liquid: tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Biguanides	Lecithin, saponin, polysorbate 80	Polysorbate 80, 30 g/l + saponin, 30 g/l + lecithin, 3 g/l. Rinse liquid: tryptone, 1 g/l + NaCl, 9 g/l; polysorbate 80, 5 g/l.
Metallic salts (Cu, Zn, Hg), organo-mercuric compounds	Sodium bisulphate, L-cysteine, sulfhydryl compounds, thioglycollic acid,	Sodium thioglycollate, 0,5 g/l or 5 g/l. L-cysteine, 0,8 g/l or 1,5 g/l. D/E neutralizing broth <sup>a</sup> Rinse liquid: sodium thioglycollate, 0,5 g/l.
NOTE See References [8] and [11].		
<sup>a</sup> D/E neutralizing broth (Dey/Engley neutralizing broth) see Annex A.		



## Bibliography

- [1] COLIPA. *Guidelines on Microbial Quality Management*, published by the European Cosmetic, Toiletry and Perfumery Association, 1997
- [2] CTFA. *Microbiology Guidelines*, published by the Cosmetic, Toiletry and Fragrance Association ISBN 1-882621-32-8, 2007
- [3] E P. *Microbiological Examination of non-sterile products*, 4th edition, published by the European Pharmacopoeia, 2002
- [4] FDA. *Bacteriological Analytical Manual*, 8th edition, published by the U.S. Food and Drug Administration, 1995, <http://www.cfsan.fda.gov/~ebam/bam-23.html>
- [5] JP 14, *General Tests — Microbial Limit test*, published by the Japanese Pharmacopoeia, 2001
- [6] USP 28, *Microbial Limit test (61)*, published by the U.S. Pharmacopoeia, 2005
- [7] ATLAS R.M. *Handbook of Microbiological Media*. CRC Press, 1993
- [8] SINGER S. The Use of Preservative Neutralizers in Diluents and Plating Media. *Cosmetics and Toiletries*. 1987 December, **102** p. 55
- [9] ISO 21149, *Cosmetics — Microbiology — Enumeration and detection of aerobic mesophilic bacteria*
- [10] ISO 18415, *Cosmetics — Microbiology — Detection of specified and non-specified microorganisms*
- [11] EN 1040, *Chemical disinfectants and antiseptics — Basic bactericidal activity — Test method and requirements (phase 1)*
- [12] ISO 29621, *Cosmetics — Microbiology — Guidelines for the risk assessment and identification of microbiologically low-risk products*



## **SLS CERTIFICATION MARK**

*The Sri Lanka Standards Institution is the owner of the registered certification mark shown below. Beneath the mark, the number of the Sri Lanka Standard relevant to the product is indicated. This mark may be used only by those who have obtained permits under the SLS certification marks scheme. The presence of this mark on or in relation to a product conveys the assurance that they have been produced to comply with the requirements of the relevant Sri Lanka Standard under a well designed system of quality control inspection and testing operated by the manufacturer and supervised by the SLSI which includes surveillance inspection of the factory, testing of both factory and market samples.*

*Further particulars of the terms and conditions of the permit may be obtained from the Sri Lanka Standards Institution, 17, Victoria Place, Elvitigala Mawatha, Colombo 08.*



## SRI LANKA STANDARDS INSTITUTION

The Sri Lanka Standards Institution (SLSI) is the National Standards Organization of Sri Lanka established under the Sri Lanka Standards Institution Act No. 6 of 1984 which repealed and replaced the Bureau of Ceylon Standards Act No. 38 of 1964. The Institution functions under the Ministry of Science, Technology and Research.

The Principal objects of the Institution as set out in the Act are to prepare standards and promote their adoption, to provide facilities for examination and testing of products, to operate a Certification Marks Scheme, to certify the quality of products meant for local consumption or exports and to promote Standardization and quality control by educational, consultancy and research and research activity.

The Institution is financed by Government grants, and by the income from the sale of its publications and other services offered for Industry and Business Sector. Financial and Administrative control is vested in a Council appointed in accordance with the provisions of the Act.

The development and formulation of National Standards is carried out by Technical Experts and representatives of other interest groups, assisted by the permanent officers of the Institution. These Technical Committees are appointed under the purview of the Sectoral Committees which in return are appointed by the Council. The Sectoral Committees give the final Technical approval for the Draft National Standards prior to the approval by the Council of the SLSI.

All members of the Technical and Sectoral Committees render their services in an honorary capacity. In this process the Institution Endeavours to ensure adequate representation of all view points.

In the International field the Institution represents Sri Lanka in the International Organization for Standardization (ISO), and participates in such fields of Standardization as are of special interest to Sri Lanka.