

SRI LANKA STANDARD 735 : PART 15:2017
(ISO 13580:2005)
UDC 579.6

METHODS OF
TESTS FOR MILK AND MILK PRODUCTS
PART 15: DETERMINATION OF TOTAL SOLID
CONTENT IN YOGHURT (REFERENCE METHOD)

SRI LANKA STANDARDS INSTITUTION

Sri Lanka Standard
METHODS OF TESTS FOR MILK AND MILK PRODUCTS
PART 15: DETERMINATION OF TOTAL SOLID CONTENT IN YOGHURT
(REFERENCE METHOD)

SLS 735 Part 15:2017
(ISO 13580: 2005)

Gr. D

Copyright Reserved
SRI LANKA STANDARDS INSTITUTION
17, Victoria Place
Elvitigala Mawatha
Colombo - 08
Sri Lanka.

Sri Lanka Standards are subject to periodical revision in order to accommodate the progress made by industry. Suggestions for improvement will be recorded and brought to the notice of the Committees to which the revisions are entrusted.

This standard does not purport to include all the necessary provisions of a contract.

© **SLSI 2017**

All right reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the SLSI.

Sri Lanka Standard
METHODS OF TESTS FOR MILK AND MILK PRODUCTS
PART 15: DETERMINATION OF TOTAL SOLID CONTENT IN YOGHURT
(REFERENCE METHOD)

NATIONAL FOREWORD

This Sri Lanka Standard was approved by the Sectoral Committee on Food Products and was authorized for adoption and publication as a Sri Lanka Standard by the Council of the Sri Lanka Standards Institution on 2017-05-04.

The Standard prescribes the test methods for determining whether the material conforms to the requirements of the relevant individual Standards and thus form a necessary adjunct to series of Sri Lanka Standard Methods of tests of milk and milk products. However, keeping in view the experience gained during the years and various International Standards brought out by the International Organization for Standardization (ISO) on the subject of testing of milk and milk products, it was decided to publish it with a view of updating the existing methods of test and by incorporating those not covered earlier.

In order to accommodate large number of test methods within the scope of one standard, this Standard is published in several Parts.

This Part of the Standard is identical with **ISO 13580 : 2005 (IDF 151)** Yoghurt - Determination of total solids content (reference method) content published by International Organization for Standardization (ISO).

TERMINOLOGY AND CONVENTIONS

The text of the International Standard has been accepted as a suitable for publication, without deviation, as a Sri Lanka Standard. However, certain terminology and conventions are not identical with those used in Sri Lanka Standards. Attention is therefore drawn to the following:

- a) Wherever the words “International Standard” appear referring to this Standard, they should be interpreted as “Sri Lanka Standard”.
- b) The comma has been used throughout as a decimal marker. In Sri Lanka Standards it is the current practice to use the full point at the base line as the decimal marker.
- c) Wherever page numbers are quoted, they are ISO page numbers.

Cross Reference

International Standards

Corresponding Sri Lanka Standard

ISO 11869 Yoghurt –Determination of titratable acidity—Potentiometric method	No corresponding Sri Lanka Standard
---	-------------------------------------

INTERNATIONAL
STANDARD

SLS 735-15:2017

ISO
13580

IDF
151

First edition
2005-05-01

**Yogurt — Determination of total solids
content (Reference method)**

*Yaourt — Détermination de la teneur totale en matières solides
(Méthode de référence)*



Reference numbers
ISO 13580:2005(E)
IDF 151:2005(E)

© ISO and IDF 2005

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. Neither the ISO Central Secretariat nor the IDF accepts any liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies and IDF national committees. In the unlikely event that a problem relating to it is found, please inform the ISO Central Secretariat at the address given below.

© ISO and IDF 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO or IDF at the respective address below.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

International Dairy Federation
Diamant Building • Boulevard Auguste Reyers 80 • B-1030 Brussels
Tel. + 32 2 733 98 88
Fax + 32 2 733 04 13
E-mail info@fil-idf.org
Web www.fil-idf.org

Published in Switzerland

Contents

Page

Foreword.....	iv
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	1
5 Reagents	1
6 Apparatus	2
7 Sampling	2
8 Preparation of test sample	2
8.1 Natural yogurt and flavoured, sweetened yogurt	2
8.2 Fruit yogurt	2
9 Procedure	3
9.1 Preparation of the dish	3
9.2 Test portion	3
9.3 Determination of total solids content	3
9.4 Determination of lactic acid content	3
10 Calculation and expression of results	4
10.1 Calculation	4
10.2 Expression of results	4
11 Precision	4
11.1 Interlaboratory test	4
11.2 Repeatability	4
11.3 Reproducibility	4
12 Test report	5
Annex A (informative) Interlaboratory test	6
Bibliography	7

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 13580|IDF 151 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF), in collaboration with AOAC International. It is being published jointly by ISO and IDF and separately by AOAC International.

Foreword

IDF (the International Dairy Federation) is a worldwide federation of the dairy sector with a National Committee in every member country. Every National Committee has the right to be represented on the IDF Standing Committees carrying out the technical work. IDF collaborates with ISO and AOAC International in the development of standard methods of analysis and sampling for milk and milk products.

Draft International Standards adopted by the Action Teams and Standing Committees are circulated to the National Committees for voting. Publication as an International Standard requires approval by at least 50 % of the National Committees casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. IDF shall not be held responsible for identifying any or all such patent rights.

ISO 13580|IDF 151 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF), in collaboration with AOAC International. It is being published jointly by ISO and IDF and separately by AOAC International.

All work was carried out by the Joint ISO/IDF/AOAC Action Team on *Water*, of the Standing Committee *Main components of milk*, under the aegis of its project leader, Mrs M. Nicolas (FR).

This edition of ISO 13580|IDF 151 cancels and replaces the first edition of IDF 151:1991.

Yogurt — Determination of total solids content (Reference method)

1 Scope

This International Standard specifies a reference method for the determination of the total solids content of plain, flavoured, sweetened and fruit yogurts.

2 Normative references

The following referenced document is indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 11869, *Yogurt — Determination of titratable acidity — Potentiometric method*¹⁾

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

total solids content

mass fraction of substances remaining after completion of the heating process specified in this International Standard

NOTE The total solids content is expressed as a percentage by mass.

4 Principle

Water from a test portion is evaporated in the presence of zinc oxide in a drying oven at $102\text{ °C} \pm 2\text{ °C}$. The lactic acid content is determined in order to compensate for the loss of water produced by neutralization.

5 Reagents

Use only reagents of recognized analytical grade, unless otherwise specified, and distilled water or demineralized water or water of equivalent purity.

5.1 Zinc oxide, at least 99 % pure.

1) Equivalent to IDF 150.

6 Apparatus

Usual laboratory equipment and, in particular, the following.

- 6.1 **Analytical balance**, capable of weighing to the nearest 1 mg, with a readability of 0,1 mg.
- 6.2 **Desiccator**, containing an effective desiccant (e.g. freshly dried silica gel with hygrometric indicator).
- 6.3 **Drying oven**, well ventilated, capable of being maintained at $102\text{ °C} \pm 2\text{ °C}$ throughout the working space.
- 6.4 **Flat-bottom dishes**, of height 20 mm to 25 mm, of diameter 50 mm to 75 mm, made of appropriate material (e.g. stainless steel, nickel or aluminium), provided with well-fitting, readily removable lids.
- 6.5 **Boiling water bath** or **steam bath**.
- 6.6 **Short stirring rods**, made of glass, flattened at one end and of suitable size to fit into the dish (6.4).
- 6.7 **Homogenizer**, for homogenizing fruit yogurt (see 8.2).
- 6.8 **Spoon** or **spatula**.

7 Sampling

A representative sample should have been sent to the laboratory. It should not have been damaged or changed during transport or storage.

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 707 | IDF 50.

Store the sample in such a way that deterioration and change in composition are prevented.

8 Preparation of test sample

8.1 Natural yogurt and flavoured, sweetened yogurt

Bring the sample to a temperature of between 20 °C and 25 °C. Mix the sample carefully by means of a spoon or spatula, using a rotary motion which passes from the lower layers to the surface layers of the sample so as to displace and mix them well.

8.2 Fruit yogurt

Bring the sample to a temperature of between 20 °C and 25 °C. Homogenize it using an appropriate device, in order to facilitate the grinding and dispersion of the fruit.

Transfer the test sample to an airtight container until the analysis, which should be carried out as soon as possible after homogenization.

If delay cannot be avoided, take all necessary precautions to prevent deterioration of the sample and to avoid condensation of moisture on the internal surface of the sample container.

9 Procedure

9.1 Preparation of the dish

Heat an open dish (6.4) containing approximately 2 g of zinc oxide (5.1), together with its lid and a stirring rod (6.6) placed on top of the lid, in the drying oven (6.3) set at 102 °C for at least 1 h.

Place the lid with the stirring rod on top on the dish and immediately transfer it to the desiccator (6.2). Allow it to cool to room temperature for at least 45 min.

Weigh the dish, with the lid and rod, to the nearest 1 mg.

9.2 Test portion

Move the zinc oxide to one side of the prepared dish (9.1) by tilting. Place on the clear space approximately 1 g of the prepared sample (Clause 8). Replace the lid on the dish with the stirring rod on top.

Weigh the dish, with the stirring rod on top, to the nearest 1 mg.

9.3 Determination of total solids content

9.3.1 Add 5 ml of water to the test portion in the dish (9.2). Using the stirring rod, thoroughly mix the diluted test portion and the zinc oxide. Spread the mixture evenly over the bottom of the dish. Leave the stirring rod with the flattened end used for mixing resting in the mixture and the other end resting on the rim of the dish.

9.3.2 Heat the dish on the boiling water bath or steam bath (6.5), ensuring that the maximum possible area of the dish is exposed to the steam. Continue the heating process for approximately 30 min, with frequent mixing of the contents of the dish during the early drying stage, so as to obtain maximum liquid evaporation.

9.3.3 Remove the dish from the water bath or steam bath and wipe its base to remove any water. Leave the stirring rod in the dish and then place it, together with the lid by its side, in the drying oven (6.3) set at 102 °C for 3 h. Then cover the dish with its lid and immediately transfer to the desiccator (6.2).

9.3.4 Allow the dish and contents to cool in the desiccator to the temperature of the weighing room for at least 45 min. Weigh the dish and contents to the nearest 1 mg.

9.3.5 Again, heat the dish and its contents, together with its lid, for a further 1 h, as described in 9.3.3.

Cover the dish with its lid and immediately transfer it to the desiccator (6.2). Allow to cool as in 9.3.4. Weigh the dish and its lid, to the nearest 1 mg.

9.3.6 Repeat this heating and weighing procedure (9.3.5) until the mass of the dish with its lid decreases by 1,0 mg or less, or increases between two successive weighings.

Take the lowest mass recorded for the calculation.

9.4 Determination of lactic acid content

In order to compensate for the water loss as a result of neutralizing the yogurt by means of zinc oxide, determine, in accordance with ISO 11869, the titratable acidity of the test sample (expressed as grams of lactic acid per 100 g of product).

10 Calculation and expression of results

10.1 Calculation

Calculate the total solids content of the test sample, w , expressed as a percentage by mass, by using the following equation:

$$w = \frac{(m_2 - m_0)}{(m_1 - m_0)} \times 100 + (0,1 a)$$

where

m_0 is the mass of the dish (including zinc oxide), lid and stirring rod (9.1), in grams;

m_1 is the mass of the dish (including zinc oxide), lid, stirring rod and test portion (9.2), in grams;

m_2 is the mass of the dish, lid, stirring rod and dried test portion (including zinc oxide) (9.3.6), in grams;

a is the titratable acidity obtained in 9.4, expressed as grams of lactic acid per 100 g of product;

0,1 is the correction value to compensate for the loss of water as a result of neutralizing the acidity of yogurt with zinc oxide (9.3.1).

10.2 Expression of results

Report the test result to two decimal places.

11 Precision

11.1 Interlaboratory test

Details of an interlaboratory test on the precision of the method are summarized in Annex A. The values derived from this interlaboratory test may not be applicable to concentration ranges and matrices other than those given.

11.2 Repeatability

The absolute difference between two independent single test results, obtained with the same method on identical test material in the same laboratory by the same operator using the same equipment within a short interval of time, will in not more than 5 % of cases be greater than 0,28 %.

11.3 Reproducibility

The absolute difference between two single test results, obtained with the same method on identical test material in different laboratories by different operators using different equipment, will in not more than 5 % of cases be greater than 0,45 %.

12 Test report

The test report shall specify:

- a) all information necessary for the complete identification of the sample;
- b) the sampling method used, if known;
- c) the test method used with reference to this International Standard;
- d) all operating details not specified in this International Standard, or regarded as optional, together with details of any incidents which may have influenced the test results;
- e) the test results obtained and, if the repeatability has been checked, the final quoted result obtained.

Annex A (informative)

Interlaboratory test

An international collaborative test (see Reference [4]) involving eight laboratories was carried out on eight yogurts. The Swiss Dairy Research Institute organized the test.

The results obtained were subjected to statistical analysis in accordance with ISO 5725-1 and ISO 5725-2 to give the precision data shown in Table A.1. All values in Table A.1 are expressed as a percentage by mass.

Table A.1 — Result of interlaboratory test

	Test samples								Mean
	A	B	C	D	E	F	G	H	
No. of participants after eliminating outliers	8	8	8	8	8	8	8	8	
Mean value, %	12,88	19,42	24,53	21,62	11,37	11,65	13,63	21,9	
Repeatability standard deviation, s_r , %	0,096	0,092	0,068	0,107	0,096	0,132	0,071	0,107	0,096
Coefficient of variation of repeatability, %	0,75	0,48	0,40	0,50	0,84	1,14	0,52	0,51	0,64
Repeatability limit r ($= 2,8 s_r$), %	0,27	0,26	0,27	0,30	0,27	0,37	0,20	0,30	0,28
Reproducibility standard deviation, s_R , %	0,150	0,196	0,207	0,182	0,132	0,157	0,093	0,175	0,16
Coefficient of variation of reproducibility, %	1,16	1,02	0,85	0,85	1,16	1,34	0,69	0,83	0,99
Reproducibility limit R ($= 2,8 s_R$), %	0,42	0,55	0,58	0,51	0,37	0,44	0,26	0,49	0,45

Bibliography

- [1] ISO 707|IDF 50, *Milk and milk products — Guidance on sampling*
- [2] ISO 5725-1, *Accuracy (trueness and precision) of measurement methods and results — Part 1: General principles and definitions*
- [3] ISO 5725-2, *Accuracy (trueness and precision) of measurement methods and results — Part 2: Basic method for the determination of repeatability and reproducibility of a standard measurement method*
- [4] *Bulletin of the International Dairy Federation*, No. 285, 1993, p. 30

SLS 735-15:2017
ISO 13580:2005(E)
IDF 151:2005(E)

ICS 67.100.99

Price based on 7 pages

SLS CERTIFICATION MARK

The Sri Lanka Standards Institution is the owner of the registered certification mark shown below. Beneath the mark, the number of the Sri Lanka Standard relevant to the product is indicated. This mark may be used only by those who have obtained permits under the SLS certification marks scheme. The presence of this mark on or in relation to a product conveys the assurance that they have been produced to comply with the requirements of the relevant Sri Lanka Standard under a well designed system of quality control inspection and testing operated by the manufacturer and supervised by the SLSI which includes surveillance inspection of the factory, testing of both factory and market samples.

Further particulars of the terms and conditions of the permit may be obtained from the Sri Lanka Standards Institution, 17, Victoria Place, Elvitigala Mawatha, Colombo 08.



SRI LANKA STANDARDS INSTITUTION

The Sri Lanka Standards Institution (SLSI) is the National Standards Organization of Sri Lanka established under the Sri Lanka Standards Institution Act No. 6 of 1984 which repealed and replaced the Bureau of Ceylon Standards Act No. 38 of 1964. The Institution functions under the Ministry of Science, Technology and Research.

The Principal objects of the Institution as set out in the Act are to prepare standards and promote their adoption, to provide facilities for examination and testing of products, to operate a Certification Marks Scheme, to certify the quality of products meant for local consumption or exports and to promote Standardization and quality control by educational, consultancy and research and research activity.

The Institution is financed by Government grants, and by the income from the sale of its publications and other services offered for Industry and Business Sector. Financial and Administrative control is vested in a Council appointed in accordance with the provisions of the Act.

The development and formulation of National Standards is carried out by Technical Experts and representatives of other interest groups, assisted by the permanent officers of the Institution. These Technical Committees are appointed under the purview of the Sectoral Committees which in return are appointed by the Council. The Sectoral Committees give the final Technical approval for the Draft National Standards prior to the approval by the Council of the SLSI.

All members of the Technical and Sectoral Committees render their services in an honorary capacity. In this process the Institution Endeavours to ensure adequate representation of all view points.

In the International field the Institution represents Sri Lanka in the International Organization for Standardization (ISO), and participates in such fields of Standardization as are of special interest to Sri Lanka.