

**SRI LANKA STANDARD 393: PART 3: 2017**  
**(ISO 6887-3: 2017)**  
**UDC 579.67**

**CODE OF PRACTICE FOR PREPARATION OF  
TEST SAMPLES, INITIAL SUSPENSION AND  
DECIMAL DILUTIONS FOR  
MICROBIOLOGICAL EXAMINATION OF  
FOOD AND ANIMAL FEEDING STUFFS  
PART 3: SPECIFIC RULES FOR THE  
PREPARATION OF FISH AND FISHERY  
PRODUCTS  
(SECOND REVISION)**

**SRI LANKA STANDARDS INSTITUTION**



**Sri Lanka Standard**  
**CODE OF PRACTICE FOR PREPARATION OF TEST SAMPLES, INITIAL  
SUSPENSION AND DECIMAL DILUTIONS FOR MICROBIOLOGICAL  
EXAMINATION OF FOOD AND ANIMAL FEEDING STUFFS**  
**Part 3: Specific Rules for the Preparation of Fish and Fishery Products**  
*(Second Revision)*

**SLS 393: PART 3: 2017**  
**(ISO 6887-3: 2017)**

**Gr. E**

*Copyright Reserved*  
**SRI LANKA STANDARDS INSTITUTION**  
**No. 17, Victoria Place**  
**Elvitigala Mawatha**  
**Colombo 08**  
**SRI LANKA**

Sri Lanka Standards are subject to periodical revision in order to accommodate the progress made by industry. Suggestions for improvement will be recorded and brought to the notice of the Committees to which the revisions are entrusted.

This Standard does not purport to include all the necessary provisions of a contract.

© SLSI 2017

All right reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the SLSI.

**Sri Lanka Standard**  
**CODE OF PRACTICE FOR PREPARATION OF TEST SAMPLES, INITIAL  
SUSPENSION AND DECIMAL DILUTIONS FOR MICROBIOLOGICAL  
EXAMINATION OF FOOD AND ANIMAL FEEDING STUFFS**  
**Part 3: Specific Rules for the Preparation of Fish and Fishery Products**  
*(Second Revision)*

**NATIONAL FOREWORD**

This Sri Lanka Standard was approved by the Sectoral Committee on Food Products and was authorized for adoption and publication as a Sri Lanka Standard by the Council of the Sri Lanka Standards Institution on 2017-07-21.

This Standard was first published in 1976 with a view of providing a general guidance with regard to the practice to be followed, precautions to be observed in the sampling of different types of foods and in the handling of the samples for microbiological analysis. However, keeping in view the experience gained during the years and various International Standards brought out by the International Organization for Standardization (ISO) on the subject of microbiology of food and animal feeding stuffs, it was decided to revise it with a view to updating the existing rules and by incorporating those not covered earlier.

For different products of food and animal feeding stuffs, it is necessary to take special precautions and in order to accommodate these precautions within the scope of one Standard, this revised Standard is published in several parts.

This part of the Standard is identical with **ISO 6887-3 : 2017**, Microbiology of the food chain – Preparation of test samples, initial suspension and decimal dilutions for microbiological examination, Part 3 – Specific rules for the preparation of fish and fishery products published by the International Organization for Standardization (ISO).

**Terminology and Conventions:**

The text of the International Standard has been accepted as suitable for publication, without deviation, as a Sri Lanka Standard. However, certain terminology and conventions are not identical with those used in Sri Lanka Standards. Attention is therefore drawn to the following:

- a) Wherever the words “International Standard” appear referring to this Standard should be interpreted as “Sri Lanka Standard”.
- b) The comma has been used throughout as a decimal marker. In Sri Lanka Standards it is the current practice to use the full point on the base line as the decimal marker.
- c) Wherever page numbers are quoted, they are ISO page numbers.

## **Cross References**

### **International Standard**

ISO 6887-1 Microbiology of the food chain-Part 1: General rules for the preparation of the initial suspension and decimal dilutions

ISO 7218: Microbiology of food and animal feeding stuffs-General requirements and guidance for microbiological examinations

### **Corresponding Sri Lanka Standard**

SLS 393 Code of practice for -preparation of test samples, initial suspension and decimal dilutions for microbiological examination of food and animal feeding stuff Part 1: General rules for the preparation of the initial suspension and decimal dilutions

SLS 1463: General requirements and guidance for microbiological examinations of food and animal feeding stuffs

---

---

**Microbiology of the food chain —  
Preparation of test samples, initial  
suspension and decimal dilutions for  
microbiological examination —**

Part 3:

**Specific rules for the preparation of  
fish and fishery products**

*Microbiologie de la chaîne alimentaire — Préparation des  
échantillons, de la suspension mère et des dilutions décimales en vue  
de l'examen microbiologique —*

*Partie 3: Règles spécifiques pour la préparation des produits de la pêche*





**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2017, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Ch. de Blandonnet 8 • CP 401  
CH-1214 Vernier, Geneva, Switzerland  
Tel. +41 22 749 01 11  
Fax +41 22 749 09 47  
copyright@iso.org  
www.iso.org



# Contents

Page

<b>Foreword</b> .....	<b>v</b>
<b>1 Scope</b> .....	<b>1</b>
<b>2 Normative references</b> .....	<b>2</b>
<b>3 Terms and definitions</b> .....	<b>2</b>
<b>4 Principle</b> .....	<b>2</b>
<b>5 Diluents</b> .....	<b>2</b>
<b>6 Apparatus</b> .....	<b>2</b>
<b>7 Sampling and sample types</b> .....	<b>3</b>
7.1 General procedures.....	3
7.2 Specific procedures for sampling bivalve molluscs, echinoderms and tunicates from primary production.....	3
7.2.1 General.....	3
7.2.2 Sampling and laboratory sample transport.....	3
7.2.3 Sampling method.....	3
7.2.4 Size and number of individuals per sample.....	4
7.2.5 Temperature control during transport.....	4
7.3 Specific procedures for sampling bivalve molluscs, gastropods, echinoderms and tunicates placed on the market.....	4
<b>8 General procedures</b> .....	<b>4</b>
<b>9 Specific procedures</b> .....	<b>4</b>
9.1 Raw fishery products, including fish, crustaceans, molluscs, tunicates and echinoderms (see <a href="#">Annex A</a> ).....	4
9.1.1 Whole fresh fish (more than 15 cm in length).....	4
9.1.2 Whole fresh fish (less than 15 cm in length).....	5
9.1.3 Sliced fish, fillets and steaks.....	5
9.1.4 Whole and sliced cephalopods.....	5
9.1.5 Whole crustacea such as crabs.....	5
9.1.6 Shelled crustacea flesh.....	5
9.1.7 Crustacea such as prawns, crayfish, and lobsters.....	6
9.1.8 Live bivalve molluscs.....	6
9.1.9 Echinoderms.....	7
9.2 Processed products.....	7
9.2.1 Whole smoked fish.....	7
9.2.2 Smoked fish fillets and slices, with or without skin.....	8
9.2.3 Whole cooked molluscs in the shell.....	8
9.2.4 Fish and fish-based multi-component products (e.g. pre-prepared fish taco, mixed seafood selections, mixed fish ball).....	8
9.2.5 Cooked or precooked shelled bivalves.....	8
9.2.6 Salted or pickled products (including fish eggs/roe such as caviar).....	8
9.2.7 Dried fish including dried salted fish.....	9
9.2.8 Fermented products.....	9
9.2.9 Marinated products.....	9
9.2.10 Breaded products.....	9
9.3 Frozen fish, crustacea, molluscs, tunicates, and echinoderms.....	9
9.3.1 Fish fillets, large fish pieces frozen in blocks, frozen small parts and single portions.....	9
9.3.2 Shelled crustacea (such as prawns) frozen in blocks.....	9
9.3.3 Whole crustacea (such as prawns) frozen in blocks.....	9
9.3.4 Flaked crustacean flesh (such as crab meats) frozen in blocks.....	9
9.3.5 Molluscs (whole cephalopods, bivalve molluscs and gastropods).....	10
<b>10 Further dilutions</b> .....	<b>10</b>

<b>Annex A (informative) Classification of major taxa</b> .....	<b>11</b>
<b>Annex B (informative) Recommended number of individual live bivalve molluscs to be submitted to the laboratory</b> .....	<b>12</b>
<b>Annex C (informative) Additional guidance for small fish, crabs and lobsters</b> .....	<b>13</b>
<b>Bibliography</b> .....	<b>16</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: [www.iso.org/iso/foreword.html](http://www.iso.org/iso/foreword.html).

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*.

This second edition cancels and replaces the first edition (ISO 6887-3:2003), which has been technically revised.

A list of all parts in the ISO 6887 series can be found on the ISO website.



# Microbiology of the food chain — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination —

## Part 3: Specific rules for the preparation of fish and fishery products

**WARNING** — The use of this document may involve hazardous materials, operations and equipment. It is the responsibility of the user of this document to establish appropriate safety and health practices and to determine the applicability of regulatory limitations before use.

### 1 Scope

This document specifies rules for the preparation of fish and fishery product samples and their suspension for microbiological examination when the samples require a different preparation from the methods described in ISO 6887-1. ISO 6887-1 defines the general rules for the preparation of the initial suspension and dilutions for microbiological examination.

This document includes special procedures for sampling raw molluscs, tunicates and echinoderms from primary production areas.

**NOTE 1** Sampling of raw molluscs, tunicates and echinoderms from primary production areas is included in this document, rather than ISO 13307, which specifies rules for sampling from the terrestrial primary production stage.

This document excludes preparation of samples for both enumeration and detection test methods where preparation details are specified in the relevant International Standards (e.g. ISO/TS 15216-1 and ISO/TS 15216-2 for determination of hepatitis A virus and norovirus in food using real-time RT-PCR).

This document is intended to be used in conjunction with ISO 6887-1. It is applicable to the following raw, processed or frozen fish and shellfish and their products (see [Annex A](#) for classification of major taxa):

- a) Raw fishery products, molluscs, tunicates and echinoderms including:
  - whole fish or fillets, with or without skin and heads, and gutted;
  - crustaceans, whole or shelled;
  - cephalopods;
  - bivalve molluscs;
  - gastropods;
  - tunicates and echinoderms.
- b) Processed products including:
  - smoked fish, whole or prepared fillets, with or without skin;
  - cooked or partially cooked, whole or shelled crustaceans, molluscs, tunicates and echinoderms;
  - cooked or partially cooked fish and fish-based multi-component products.

- c) Raw or cooked frozen fish, crustaceans, molluscs and others, in blocks or otherwise, including:
- fish, fish fillets and pieces;
  - whole and shelled crustacean (e.g. flaked crab, prawns), molluscs, tunicates and echinoderms.

NOTE 2 The purpose of examinations performed on these samples can be either hygiene testing or quality control. However, the sampling techniques described in this document relate mainly to hygiene testing (on muscle tissues).

## 2 Normative references

The following documents are referred to in text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 6887-1, *Microbiology of the food chain — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination — Part 1: General rules for the preparation of the initial suspension and decimal dilutions*

ISO 7218, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations*

## 3 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 6887-1 apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

## 4 Principle

The general principles for sample preparation and subsequent steps are described in ISO 6887-1. This document describes specific measures for fish and fishery products, including raw, processed and frozen products.

## 5 Diluents

Diluents for general use and special purposes are described in ISO 6887-1 and there are no additional specific requirements for fish and fishery products.

## 6 Apparatus

Usual microbiological laboratory equipment for general use (ISO 7218 and ISO 6887-1) and in particular, the following:

### 6.1 Homogenizer.

**6.1.1 Rotary homogenizer** (blender), as specified in ISO 7218, but if a large test portion is used, the equipment should include a 1 l bowl.

**6.1.2 Peristaltic homogenizer**, as specified in ISO 7218.

**6.2 Sterile instruments** suitable for dissecting samples and opening shells (e.g. oyster knives, hammer, pliers, adjustable vice, oyster cracker, sterile scissors, shellfish picker, winkle picker, scalpels and butcher's knives).

**6.3 Sterile forceps** (small and large), **spatulas** and **spoons**.

**6.4 Small stiff brush**, for scrubbing shells.

**6.5 Electric drill**, equipped with sterile wood bit (14 mm or 16 mm diameter).

**6.6 Sterile gauze sheets**, suitable for preventing splintering when breaking up shells.

**6.7 Food grade plastic bags with waterproof labels**, suitable as sampling containers.

**6.8 Gloves**, strong, suitable to protect operator from injury.

## **7 Sampling and sample types**

### **7.1 General procedures**

Carry out sampling in accordance with the instructions given in this clause for samples from the primary production stage (7.2) or products placed on the market (7.3). For products not detailed here, carry out sampling according to the specific standard appropriate to the product concerned or see ISO/TS 17728. If specific sampling instructions are not available, it is recommended that agreement be reached on this subject by the parties concerned.

### **7.2 Specific procedures for sampling bivalve molluscs, echinoderms and tunicates from primary production**

#### **7.2.1 General**

The design and implementation of environmental sampling programmes will affect the results obtained from microbiological examinations. Where the results of this testing are used in microbiological monitoring programmes, particularly for official controls such as classification and monitoring of marine production areas, special consideration should be given to formal recording of sampling plans, species selection and spatial and temporal aspects of sampling design.<sup>[6]</sup>

#### **7.2.2 Sampling and laboratory sample transport**

A sampling protocol containing details of sampling methods, cleaning, packing and transport requirements should be agreed by the parties concerned.

#### **7.2.3 Sampling method**

The species under examination should be sampled, as far as possible, using the method employed for commercial harvesting. Equipment used for sampling shall be restricted to that used for this purpose. To avoid contamination by micro-organisms adhering to marine sediments, disturbance of surrounding sediments shall be avoided. Once removed from water and having closed, animals shall be cleaned by rinsing or scrubbing with clean seawater or fresh potable water. Animals shall not be re-immersed in water.

Individual laboratory samples shall be placed in separate undamaged food grade plastic bags (6.7) or equivalent, with waterproof labels containing information to ensure sample traceability.

Where sampling using the commercial harvesting method is not possible, unprocessed animals harvested for commercial purpose should be taken periodically as checks to show that results for laboratory samples collected using the alternative sampling method are acceptable.

#### 7.2.4 Size and number of individuals per sample

Laboratory samples should comprise individuals within the normal commercial size range. A pooled sample comprising a minimum of 10 animals with a minimum amount of flesh and intravalvular liquid of 50 g should be used (for very small species such as *Donax* spp. a minimum amount of 25 g is permitted). Additional animals shall be collected to allow for a proportion of individuals received at the laboratory being in a moribund state. Recommended numbers of individuals for some species are given in [Annex B](#).

#### 7.2.5 Temperature control during transport

The temperature of the sample (either the laboratory sample or the surrounding seawater) should be recorded immediately after collection.

Transport temperature shall be between 0 °C and 10 °C and the equipment used shall be capable of achieving this temperature range within 4 h of sample packing and maintaining it for at least 24 h. If cool packs are used, laboratory samples shall not come into direct contact with their surfaces. Samples shall not be frozen.

Internal air temperature of the temperature controlled unit shall be recorded on receipt at the laboratory.

For samples where less than 4 h have elapsed between collection from the production area and receipt at the laboratory, internal air/sample temperature should be less than the temperature recorded at the time of sampling.

Microbiological examination should be initiated within 24 h of collection of the sample from the production area. If testing cannot be initiated within 24 h or if sample temperatures between 0 °C and 10 °C cannot be achieved, data should be generated to verify that the use of alternate transport and storage conditions does not affect the microbiological content of the sample.

NOTE Studies have shown that *E. coli* will not significantly increase in mussels (*Mytilus edulis*) or Pacific oysters (*Crassostrea gigas*) at temperatures of 15 °C or less for up to 48 h.<sup>[8]</sup>

### 7.3 Specific procedures for sampling bivalve molluscs, gastropods, echinoderms and tunicates placed on the market

Use the specific sampling procedures given in [7.2.4](#).

## 8 General procedures

All preparations and manipulations shall be carried out using aseptic techniques and sterile equipment (ISO 7218).

## 9 Specific procedures

### 9.1 Raw fishery products, including fish, crustaceans, molluscs, tunicates and echinoderms (see [Annex A](#))

#### 9.1.1 Whole fresh fish (more than 15 cm in length)

The gills and the anus shall be covered with sterile cotton wool, drenched in alcohol at a volume fraction of 70 %. Disinfect the surface of the dorsal region (using cotton wool with alcohol at a volume fraction



of 70 %) and remove and discard a section of the skin using sterile forceps (6.3) and scalpel (6.2). Take a cube-shaped sample of dorsal muscle, dice it and break up in an appropriate diluent.

If the fish is eviscerated, the gills shall be covered with sterile cotton wool, drenched in alcohol at a volume fraction of 70 %, and a cube-shaped sample of dorsal muscle shall be removed from inside the body cavity.

Use sufficient material from the laboratory sample to give a representative test portion as specified in the test method.

Add diluent to make a 1 in 10 suspension and blend in a rotary or peristaltic homogenizer (6.1) as necessary.

#### **9.1.2 Whole fresh fish (less than 15 cm in length)**

Using sterile scissors (6.2) and forceps (6.3), remove a portion of fish just anterior to the tail insertion by making two cuts to produce transverse sections, the first cut to remove the tail and tail insertion and the second to remove a steak.

Use sufficient material from the laboratory sample to give a representative test portion as specified in the test method.

Add diluent to make a 1 in 10 suspension and blend in a rotary or peristaltic homogenizer (6.1) as necessary.

NOTE Additional guidance for small fish up to 15 cm in length is given in [Annex C](#).

#### **9.1.3 Sliced fish, fillets and steaks**

No specific requirements; treat in accordance with ISO 6887-1.

#### **9.1.4 Whole and sliced cephalopods**

Disinfect the surface of the skin and suckers (using cotton wool with alcohol at a volume fraction of 70 %). Remove the skin and suckers with sterile forceps (6.3) and a scalpel (6.3) and discard. Take cube-shaped samples of dorsal muscles and pieces from the tentacles. Use sufficient material from the laboratory sample to give a representative test portion as specified in the test method.

The flesh from cephalopods is relatively firm; grind up the test portion in diluent using a rotary homogenizer (6.1.1) or cut it into fine pieces. Add further diluent to make a 1 in 10 suspension and blend in a rotary or peristaltic homogenizer (6.1) as necessary.

#### **9.1.5 Whole crustacea such as crabs**

Disinfect the surface (using cotton wool with alcohol at a volume fraction of 70 %) and with sterile hammer (6.2), pliers (6.2) or forceps (6.3) remove or break the carapace (see C.2) and claws to extract the maximum amount of flesh for testing. For large claws, an oyster cracker (6.2) can be used to break open the shell before extracting the flesh.

Use sufficient material from the laboratory sample to give a representative test portion as specified in the test method.

Add diluent to make a 1 in 10 suspension and blend in a rotary or peristaltic homogenizer (6.1) as necessary.

#### **9.1.6 Shelled crustacea flesh**

Take the amount of flesh required in the test method, make the initial 1 in 10 suspension in a diluent and blend in a rotary or peristaltic homogenizer (6.1) as necessary.

Use sufficient material from the laboratory sample to give a representative test portion as specified in the test method.

### 9.1.7 Crustacea such as prawns, crayfish, and lobsters

#### 9.1.7.1 Species where tails only are consumed

Disinfect the surface (using cotton wool with alcohol at a volume fraction of 70 %). Break the crustacean at the junction between the cephalothorax and abdomen (see [Figure C.3](#)). Using sterile forceps ([6.3](#)) pull the edible portion of flesh from the cephalothorax and butt end of the abdomen.

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Add the necessary quantity of diluent to give a 1 in 10 suspension.

Blend in a rotary or peristaltic homogenizer ([6.1](#)) as necessary.

#### 9.1.7.2 Species consumed whole

Take the entire animal for examination. Use sufficient material from the laboratory sample to give a representative test portion as specified in the test method.

Add the necessary quantity of diluent to give a 1 in 10 suspension.

Blend in a rotary or peristaltic homogenizer ([6.1](#)) as necessary.

### 9.1.8 Live bivalve molluscs

#### 9.1.8.1 General

On arrival at the laboratory, the internal air temperature of the transport container shall be recorded. For samples where more than 4 h have elapsed between collection and receipt, the internal air temperature should be between 0 °C and 10 °C. If the internal air temperature of the transport container is greater than 10 °C, the sample temperature should be measured; this should not exceed 10 °C. For samples where less than 4 h have elapsed between collection and receipt, internal air/sample temperature should be less than the temperature recorded at the time of sampling.

Laboratory samples shall be stored at 3 °C ± 2 °C.

The animals shall be alive. Discard individuals with open or damaged shells. A representative test sample shall contain at least 10 individuals<sup>[Z]</sup> and shall be at least 50 g (25 g for small animals, e.g. *Donax* spp.) as detailed in [7.2.4](#) above. Testing of bivalves includes both the flesh and intravalvular water; open sufficient shellfish to yield the amount of flesh and intravalvular fluid specified in the test method.

Microbiological examination should be initiated within 24 h of collection of the sample. If testing cannot be initiated within 24 h or if sample temperatures of 0 °C and 10 °C cannot be achieved, data should be generated to verify that the use of alternate transport and storage conditions does not affect the microbiological content of the sample.

NOTE Studies have shown that *E. coli* will not significantly increase in mussels (*Mytilus edulis*) or Pacific oysters (*Crassostrea gigas*) at temperatures of 15 °C or less for up to 48 h.<sup>[8]</sup>

#### 9.1.8.2 Methods requiring a 1 in 10 initial suspension

Wash and brush ([6.4](#)) each shell under running water of potable quality, especially around the hinge or opening.

Drain the cleaned bivalves and put them on a clean surface.

If there is a byssus muscle, do not tear it away; cut it with sterile scissors, knife or scalpel (6.2) before fully opening.

As each shell is opened, collect the flesh and intravalvular water in a sterile container suitable for blending. Bivalves that have lost their intravalvular water may be used if they are still alive when the shell is opened.

Add one part of flesh and intravalvular water to two parts of diluent. Blend with a rotary homogenizer (6.1.1) for 30 s to 2 min depending on the homogenizer used (ISO 7218). A peristaltic homogenizer (6.1.2) may be used but note that shell splinters can puncture plastic bags. Double- or triple-bagging can help to prevent leaking and the risk of contamination.

In this way, an approximate 1 in 3 suspension is obtained to which the required amount of diluent is added to obtain an accurate 1 in 10 initial suspension.

### 9.1.8.3 Methods requiring a 1 in 2 initial suspension

Proceed as in 9.1.8.2 but use one part of flesh and intravalvular water to one part of diluent to produce an accurate initial 1 in 2 suspension.

**NOTE** An initial suspension of 1 in 2 is required for official control testing of bivalve shellfish, marine gastropods and echinoderms according to ISO 16649-2 or other applications where a level of detection of  $\leq 200$  cfu per 100 g product is required.

### 9.1.9 Echinoderms

#### 9.1.9.1 Echinoderms such as sea urchins

Wash at least 10 individuals under running potable water, and place them on a sterile tray.

Hold the sea urchin with forceps (6.3) or wear a strong clean glove (6.8) and cut off a piece of the ventral surface with sterile sharp scissors (6.2) to expose the flesh. Collect the whole flesh and fluid in a sterile container suitable for blending.

Prepare an initial suspension of approximately 1 in 3 in diluent, homogenize in a rotary or peristaltic homogenizer (6.1) as necessary and then add the required amount of diluent to obtain an accurate 1 in 10 suspension.

#### 9.1.9.2 Echinoderms such as holothurians (e.g. sea cucumbers) and tunicates

Wash at least 10 individuals under running potable water, and place them on a sterile tray.

Cut individuals into fine pieces with sterile scissors (6.2).

Prepare an initial suspension of approximately 1 in 3 in diluent, homogenize in a rotary or peristaltic homogenizer (6.1) as necessary and then add the required amount of diluent to obtain an accurate 1 in 10 suspension.

## 9.2 Processed products

### 9.2.1 Whole smoked fish

If the whole fish is eaten, then the skin shall be included in the sample. If the skin is not eaten then the skin shall be excluded. Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

The test portion shall be taken from the dorsal area and the flesh cut, diced and homogenized using a rotary or peristaltic homogenizer (6.1) as necessary in diluent to obtain a 1 in 10 suspension.

### **9.2.2 Smoked fish fillets and slices, with or without skin**

Take pieces of the fillet and dice them, under sterile conditions, without removing the skin. Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using either a rotary or peristaltic homogenizer (6.1) as necessary in diluent to obtain a 1 in 10 suspension.

### **9.2.3 Whole cooked molluscs in the shell**

#### **9.2.3.1 Cooked or partially cooked gastropods**

Remove the operculum with a sterile scalpel (6.2) then extract the body using forceps (6.3), a winkle picker or shellfish picker (6.2).

Alternatively, carefully crush the shells open using a hammer (6.2) without damaging the flesh.

Remove any shell debris with sterile forceps (6.3) and dice the flesh.

Prepare an initial suspension of approximately 1 in 3 in diluent, homogenize, and then add the required amount of diluent to obtain an accurate 1 in 10 suspension.

#### **9.2.3.2 Cooked or partially cooked bivalves**

Extract the body from the shell using sterile forceps (6.3), scalpel and oyster knife or shellfish picker (6.2).

Dice the flesh.

Prepare an initial suspension of approximately 1 in 3 in diluent, homogenize in a rotary or peristaltic homogenizer (6.1), and then add the required amount of diluent to obtain an accurate 1 in 10 suspension.

#### **9.2.3.3 Whole cooked or partially cooked crustacea**

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Add the necessary quantity of diluent to give a 1 in 10 suspension.

Blend in a rotary or peristaltic homogenizer (6.1).

### **9.2.4 Fish and fish-based multi-component products (e.g. pre-prepared fish taco, mixed seafood selections, mixed fish ball)**

Take representative parts of each component in proportion to the amounts in the whole product. Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Add the necessary quantity of diluent to give a 1 in 10 suspension.

Blend in a rotary or peristaltic homogenizer (6.1).

### **9.2.5 Cooked or precooked shelled bivalves**

No specific requirements; treat in accordance with ISO 6887-1.

### **9.2.6 Salted or pickled products (including fish eggs/roe such as caviar)**

Treat as dehydrated or acidic products in accordance with ISO 6887-1.

### **9.2.7 Dried fish including dried salted fish**

Treat as dehydrated products in accordance with ISO 6887-1.

### **9.2.8 Fermented products**

Treat as acidic products in accordance with ISO 6887-1.

### **9.2.9 Marinated products**

Treat as acidic products in accordance with ISO 6887-1.

### **9.2.10 Breaded products**

No specific requirements; treat in accordance with ISO 6887-1.

## **9.3 Frozen fish, crustacea, molluscs, tunicates, and echinoderms**

### **9.3.1 Fish fillets, large fish pieces frozen in blocks, frozen small parts and single portions**

Either take a test portion from the frozen block using a drill with a sterile bit (6.5) or defrost at ambient temperature (18 °C to 27 °C) for approximately 60 min but no more than 3 h. Remove pieces with sterile pliers or forceps. Leave to defrost further if necessary until soft enough to cut into smaller pieces with a sterile knife (6.2) and forceps (6.3).

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Blend the pieces in a rotary or peristaltic homogenizer (6.1) with diluent to obtain a 1 in 10 suspension.

### **9.3.2 Shelled crustacea (such as prawns) frozen in blocks**

Leave the laboratory sample to defrost for approximately 60 min but no more than 3 h at ambient temperature (18 °C to 27 °C) until the block breaks. Carefully separate the block into pieces using a sterile hammer or butcher's knife (6.2) and take pieces of flesh with sterile forceps (6.3) or pliers (6.2). Alternatively remove the test portion from the frozen block using a drill with a sterile bit (6.5).

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using a rotary or peristaltic homogenizer (6.1) in diluent to obtain a 1 in 10 suspension.

### **9.3.3 Whole crustacea (such as prawns) frozen in blocks**

Leave to defrost for approximately 60 min but no more than 3 h at ambient temperature (18 °C to 27 °C) until the block breaks. Extract the individual animals with sterile pliers (6.2) or forceps (6.3). Allow to defrost so cephalothorax and abdomen (see Annex C) may be separated and the edible portion removed with sterile forceps (6.3).

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using a rotary or peristaltic homogenizer (6.1) in diluent to obtain a 1 in 10 suspension.

### **9.3.4 Flaked crustacean flesh (such as crab meats) frozen in blocks**

Remove the test portion from the frozen block using a drill with sterile bit (6.5) or defrost at ambient temperature (18 °C to 27 °C) for approximately 60 min but no more than 3 h until the block breaks. Remove pieces of flesh with sterile pliers (6.2) or forceps (6.3).

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using a rotary or peristaltic homogenizer (6.1) in diluent to obtain a 1 in 10 suspension.

### **9.3.5 Molluscs (whole cephalopods, bivalve molluscs and gastropods)**

#### **9.3.5.1 Whole cephalopods frozen in blocks**

Remove material using a drill with a sterile bit (6.5) or defrost at ambient temperature (18 °C to 27 °C) for approximately 60 min but no more than 3 h. Cut off pieces with sterile scissors or butcher's knife (6.2).

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using a rotary or peristaltic homogenizer (6.1) in diluent to obtain a 1 in 10 suspension.

#### **9.3.5.2 Whole gastropods and bivalve molluscs frozen in blocks**

Leave to defrost for approximately 60 min but no more than 3 h at ambient temperature (18 °C to 27 °C) until the block breaks. Extract the individual animals with sterile pliers (6.2) or forceps (6.3). Leave to defrost further if necessary until soft enough to extract the body from the shell using sterile forceps (6.3), scalpel and oyster knife or shellfish picker (6.2).

Alternatively crush the shells open using a sterile hammer (6.2) without damaging the flesh.

Remove any shell debris with sterile forceps (6.3) and dice the flesh.

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using a rotary or peristaltic homogenizer (6.1) in diluent to obtain a 1 in 10 suspension.

#### **9.3.5.3 Cooked or partially cooked, shelled molluscs such as gastropods and bivalve molluscs frozen in blocks**

Leave to defrost for approximately 60 min but no more than 3 h at ambient temperature (18 °C to 27 °C) until the block breaks. Extract the individual animals with sterile pliers (6.2) or forceps (6.3).

Use sufficient material from the edible portion of the laboratory sample to give a representative test portion as specified in the test method.

Homogenize using a rotary or peristaltic homogenizer (6.1) in diluent to obtain a 1 in 10 suspension.

## **10 Further dilutions**

Prepare further dilutions in accordance with ISO 6887-1.

## Annex A (informative)

### Classification of major taxa

Taxonomical Division		Examples
Phylum — Chordata	Class — Myxini	Hagfish, Nuta-unagi, Meokjango, Yu sheng
	Class — Petromyzontida	Lamprey
	Class — Chondrichthyes	Whitefish, Makorepe, ghost shark
	Class — Elasmobranchii	Sharks, flake, sora, rays, skates
	Class — Actinopterygii	Fin fish
Phylum — Arthropoda, subphylum — Crustacea		Crayfish, yabby, marron, scampi, clawed lobster, spiny lobster, langoustines, shrimp, prawns, crabs,
Phylum — Mollusca	Class — Cephalopoda	Octopus, squid, cuttlefish, nautilus.
	Class — Bivalvia	Oysters, mussels, scallops, clams, cockles
	Class — Gastropoda	Abalone (paua), conch, periwinkles, whelks, limpets, sea slugs, snails
Phylum — Chordata, subphylum — Tunicata		Sea squirts, sea pork, sea tulips, sea violet, piure
Phylum — Echinodermata	Class — Holothurian	Sea cucumber, trepan, sea slug
	Class — Echinoidea	Sea urchins (hota, ututuk, kina, uni) star fish

## Annex B (informative)

### Recommended number of individual live bivalve molluscs to be submitted to the laboratory

Species		Number
Scientific name	Common name (English)	
<i>Pecten maximus</i>	King scallop	12 to 18
<i>Aequipecten opercularis</i>	Queen scallop	18 to 35
<i>Crassostrea gigas</i>	Pacific oyster	12 to 18
<i>Ostrea edulis</i>	Flat oyster	12 to 18
<i>Mercenaria mercenaria</i>	Hard clams	12 to 18
<i>Tapes philippinarum</i>	Manilla clam	18 to 35
<i>Ruditapes decussatus</i>	Grooved carpet shells	18 to 35
<i>Spisula solida</i>	Thick trough shells	35 to 55
<i>Mya arenaria</i>	Sand gapers	12 to 18
<i>Ensis</i> spp.	Razor clams	12 to 18
<i>Mytilus</i> spp.	Mussels	18 to 35
<i>Cerastoderma edule</i>	Cockles	35 to 55
<i>Donax</i> spp.	Bean clams	40 to 70




## Annex C (informative)

### Additional guidance for small fish, crabs and lobsters

#### C.1 Small fish (up to 15 cm long)

Using sterile scissors and forceps, remove a portion of the fish just anterior to the tail insertion by making two cuts to produce transverse sections; the first cut to remove the tail and the tail insertion and the second to remove a steak (see [Figure C.1](#)).

Take care not to remove any viscera or gut contents.




#### Key

- 1 cut 1
- 2 cut 2

Figure C.1 — Example of test sampling of a fish up to 15 cm in length

## C.2 Crabs

Lift off carapace (see [Figure C.2](#)) with sterile forceps and crack claws. Using sterile forceps, take sufficient flesh to yield the amount specified in the test method.



**Key**

1 carapace


**Figure C.2 — Carapace of a crab**

### C.3 Flesh from lobsters and crayfish

Break the crustacean at the junction between the cephalothorax and abdomen (see [Figure C.3](#)).

Using sterile forceps, pull the flesh from the cephalothorax and butt end of the abdomen (this includes a little gut, which is usually eaten).

Take sufficient flesh to yield the amount specified in the test method.



**Key**

- 1 cephalothorax
- 2 abdomen

**Figure C.3 — Cephalothorax and abdomen of a lobster**

## Bibliography

- [1] ISO 13307, *Microbiology of food and animal feed — Primary production stage — Sampling techniques*
- [2] ISO/TS 15216-1, *Microbiology of the food chain — Horizontal method for determination of hepatitis A virus and norovirus in food using real-time RT-PCR — Part 1: Method for quantification*
- [3] ISO/TS 15216-2, *Microbiology of food and animal feed — Horizontal method for determination of hepatitis A virus and norovirus in food using real-time RT-PCR — Part 2: Method for qualitative detection*
- [4] ISO 16649-2, *Microbiology of food and animal feeding stuffs — Horizontal method for the enumeration of beta-glucuronidase-positive Escherichia coli — Part 2: Colony-count technique at 44 °C using 5-bromo-4-chloro-3-indolyl beta-D-glucuronide*
- [5] ISO/TS 17728, *Microbiology of the food chain — Sampling techniques for microbiological analysis of food and feed samples*
- [6] European Union Reference Laboratory for Monitoring Bacteriological and Viral Contamination of Bivalve Molluscs. Microbiological monitoring of bivalve mollusc harvesting areas, guide to good practice: Technical application ( 2014). Available at: [https://eur1cefasc.org/media/13828/gpg\\_issue-5\\_final\\_all.pdf](https://eur1cefasc.org/media/13828/gpg_issue-5_final_all.pdf)
- [7] European Communities 2005. Commission Regulation (EC) 2073/2005 on microbiological criteria for foodstuffs. *Off. J. Eur. Communities* L338, 22.12.05: 1-26
- [8] Centre for Environment Fisheries and Aquaculture Science (CEFAS). Fish Health Inspectorate (FHI) Annual Report (2015). Published: 21 May 2015. Updated: 23 February 2016. Available at: <https://www.gov.uk/government/publications/fish-health-inspectorate-reports-2015>





## **SLS CERTIFICATION MARK**

*The Sri Lanka Standards Institution is the owner of the registered certification mark shown below. Beneath the mark, the number of the Sri Lanka Standard relevant to the product is indicated. This mark may be used only by those who have obtained permits under the SLS certification marks scheme. The presence of this mark on or in relation to a product conveys the assurance that they have been produced to comply with the requirements of the relevant Sri Lanka Standard under a well designed system of quality control inspection and testing operated by the manufacturer and supervised by the SLSI which includes surveillance inspection of the factory, testing of both factory and market samples.*

*Further particulars of the terms and conditions of the permit may be obtained from the Sri Lanka Standards Institution, 17, Victoria Place, Elvitigala Mawatha, Colombo 08.*



## **SRI LANKA STANDARDS INSTITUTION**

The Sri Lanka Standards Institution (SLSI) is the National Standards Organization of Sri Lanka established under the Sri Lanka Standards Institution Act No. 6 of 1984 which repealed and replaced the Bureau of Ceylon Standards Act No. 38 of 1964. The Institution functions under the Ministry of Science & Technology.

The principal objects of the Institution as set out in the Act are to prepare standards and promote their adoption, to provide facilities for examination and testing of products, to operate a Certification Marks Scheme, to certify the quality of products meant for local consumption or exports and to promote standardization and quality control by educational, consultancy and research activity.

The Institution is financed by Government grants, and by the income from the sale of its publications and other services offered for Industry and Business Sector. Financial and administrative control is vested in a Council appointed in accordance with the provisions of the Act.

The development and formulation of National Standards is carried out by Technical Experts and representatives of other interest groups, assisted by the permanent officers of the Institution. These Technical Committees are appointed under the purview of the Sectoral Committees which in turn are appointed by the Council. The Sectoral Committees give the final Technical approval for the Draft National Standards prior to the approval by the Council of the SLSI.

All members of the Technical and Sectoral Committees render their services in an honorary capacity. In this process the Institution endeavours to ensure adequate representation of all view points.

In the International field the Institution represents Sri Lanka in the International Organization for Standardization (ISO), and participates in such fields of standardization as are of special interest to Sri Lanka.